

Procter & Gamble Co.

-PE Product Labels –

-continued



Drug Facts	
Active ingredients (in each 15 mL)	Purpose
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant
Uses	
temporarily relieves common cold symptoms: • nasal congestion • cough due to minor throat & bronchial irritation	
• helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.	
Warnings	
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema • a sodium-restricted diet	
When using this product, do not use more than directed.	
Stop use and ask a doctor if • you get nervous, dizzy or sleepless • symptoms do not improve within 7 days or occur with a fever • cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache These could be signs of a serious condition.	
If pregnant or breast-feeding , ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	

Drug Facts (continued)	
Directions	
• take only as directed • only use the dose cup provided • do not exceed 4 doses per 24 hrs	
adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	7.5 mL every 4 hrs
children under 4 yrs	do not use
Other information	
• each 15 mL contains: sodium 44 mg • Store at no greater than 25°C.	
Inactive ingredients	
anhydrous citric acid, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose	
Questions?	
1-800-362-1683	

TAMPER EVIDENT: Do not use if printed safety seal under cap is missing or damaged.
DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202
Made in Canada Patents: www.pg.com/patents
Close cap tightly  www.vicks.com

 www.pg.com
 **PEEL BACK FOR DRUG FACTS**
CLOSE TIGHTLY



**DAY & NIGHT
PACK**

children's Cough Congestion

Phenylephrine HCl - nasal decongestant
Dextromethorphan HBr - cough suppressant
Guaifenesin - expectorant

Non-Drowsy

FREE of:
Artificial Dyes & Flavors,
High Fructose Corn Syrup & Alcohol

• Cough
• Chest congestion
• Stuffy nose

Ages 4+

children's Cough Congestion Night

Phenylephrine HCl - nasal decongestant
Diphenhydramine HCl -
antihistamine/cough suppressant

Nighttime

• Cough
• Runny nose
• Stuffy nose
• Sneezing

Ages 6+

Children's Cough Congestion Day

Drug Facts (continued)

Other information
each 15 mL contains: sodium 44 mg
Store at no greater than 25°C.

Inactive ingredients anhydrous citric acid, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions? 1-800-362-1683

TAMPER EVIDENT: Do not use if printed safety seal under cap is missing or damaged.
DIST. BY: PROCTER & GAMBLE,
CINCINNATI, OH 45202
Made in Canada
Patents:
www.pg.com/
patents
www.vicks.com
P&G
www.pg.com

Children's Cough Congestion Night

Drug Facts (continued)

Directions • take only as directed
• only use the dose cup provided
• do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	do not use unless directed by a doctor
children under 4 yrs	do not use

Other information
each 15 mL contains: sodium 44 mg
Store at no greater than 25°C.

Inactive ingredients anhydrous citric acid, flavor, glycerin, polysorbate 20, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

Questions? 1-800-362-1683

2 BOTTLES - 6 FL OZ (177 mL) EACH; TOTAL 12 FL OZ (354 mL)



Children's Cough Congestion Night

Drug Facts	
Active ingredients (in each 15 mL)	Purpose
Diphenhydramine HCl 12.5 mg	Antihistamine/Cough suppressant
Phenylephrine HCl 5 mg	Nasal decongestant
Uses	
temporarily relieves common cold symptoms:	
• nasal congestion • runny nose & sneezing	
• cough due to minor throat & bronchial irritation	
Warnings	
Do not use	
• with any other product containing diphenhydramine, even one used on skin • to make a child sleepy	
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have	
• heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or occurs with smoking, asthma or emphysema • a sodium-restricted diet	
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers	
When using this product	
• do not use more than directed.	
• excitability may occur, especially in children	
• marked drowsiness may occur	
• alcohol, sedatives, and tranquilizers may increase drowsiness	
• avoid alcoholic drinks	
• be careful when driving a motor vehicle or operating machinery	
Stop use and ask a doctor if	
• you get nervous, dizzy or sleepless	
• symptoms do not improve within 7 days or occur with a fever	
• cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache	
These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	

Children's Cough Congestion Day

Drug Facts	
Active ingredients (in each 15 mL)	Purpose
Dextromethorphan HBr 10 mg	Cough suppressant
Guaifenesin 200 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant
Uses	
temporarily relieves common cold symptoms:	
• nasal congestion • cough due to minor throat & bronchial irritation	
• helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.	
Warnings	
Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have	
• heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema • a sodium-restricted diet	
When using this product, do not use more than directed.	
Stop use and ask a doctor if	
• you get nervous, dizzy or sleepless	
• symptoms do not improve within 7 days or occur with a fever	
• cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache	
These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use.	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions	
• take only as directed	
• only use the dose cup provided • do not exceed 4 doses per 24 hrs	
adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	7.5 mL every 4 hrs
children under 4 yrs	do not use



VICKS

children's
Cough Congestion Night

Phenylephrine HCl - nasal decongestant
Diphenhydramine HCl - antihistamine/cough suppressant

FREE of:
Artificial Dyes & Flavors,
High Fructose Corn Syrup & Alcohol

• Cough
• Runny nose
• Stuffy nose
• Sneezing

Ages 6+

6 FL OZ (177ml)

PARENTS:
Learn about how medicine works
www.SuppMedicineUse.org

Drug Facts

Active ingredients (in each 15 mL)
Diphenhydramine HCl 12.5 mg
Phenylephrine HCl 5 mg

Purpose
Antihistamine/Cough suppressant
Nasal decongestant

Uses
temporarily relieves common cold symptoms:
• cough due to minor throat & bronchial irritation
• nasal congestion
• runny nose & sneezing

Warnings
Do not use
• with any other product containing diphenhydramine, even one used on skin
• to make a child sleepy
• if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
• heart disease • high blood pressure • thyroid disease • diabetes
• glaucoma • trouble urinating due to enlarged prostate gland • cough that occurs with low-mach phlegm (mucus)
• a breathing problem or chronic cough that lasts or occurs with smoking, asthma or emphysema
• a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers

When using this product
• do not use more than directed.
• drowsiness may occur, especially in children
• avoid alcoholic drinks
• be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if
• you get nervous, dizzy or sleepless
• symptoms do not improve within 7 days or occur with a fever
• cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache
These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
• take only as directed
• only use the dose cup provided
• do not exceed 4 doses per 24 hrs
adults & children 12 yrs & over 30 mL every 4 hrs
children 6 to under 12 yrs 15 mL every 4 hrs
children 4 to under 6 yrs do not use unless directed by a doctor
children under 4 yrs do not use

Other information
• each 15 mL contains: sodium 44 mg
• Store at or below 25°C.

Inactive ingredients
anhydrous citric acid, flavor, glycerin, polybutene 20, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose

TAMPER EVIDENT: Do not use if printed safety seal under cap is missing or damaged.
DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202
Made in Canada
Close cap tightly
91969776

P&G
www.pg.com

Patents: www.pg.com/patents
www.vicks.com

PEEL BACK FOR DRUG FACTS

CLOSE TIGHTLY

Drug Facts (continued) If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions • take only as directed • do not exceed 8 caplets per 24 hrs adults & children 12 yrs & over 2 caplets with water every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use Other information • store at no greater than 25°C Inactive ingredients crospravodone, D&C Yellow No. 10 Aluminum Lake, FD&C Blue No. 1 Aluminum Lake, FD&C Blue No. 2 Aluminum Lake, FD&C Yellow No. 6 Aluminum Lake, flavor, magnesium stearate, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide Questions? 1-800-362-1683	Drug Facts (continued) If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms. Directions • take only as directed • do not exceed 8 caplets per 24 hrs adults & children 12 yrs & over 2 caplets with water every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use Other information • each caplet contains: sodium 4 mg • store at no greater than 25°C Inactive ingredients crospravodone sodium, crospravodone, FD&C Yellow No. 6 Aluminum Lake, flavor, maltodextrin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, silicon dioxide, stearic acid, sucralose, talc, titanium dioxide Questions? 1-800-362-1683
---	---

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

TAMPER EVIDENT: This package is safety-sealed & child resistant. Use only if blisters are intact. If difficult to open, use scissors.



ACTUAL SIZE



DayQuil
VapoCOOL[®]
SEVERE COLD & FLU
+ CONGESTION
Acetaminophen, Guaifenesin,
Phenylephrine HCl, Dextromethorphan HBr

NyQuil
VapoCOOL[®]
SEVERE COLD & FLU
+ CONGESTION
Acetaminophen, Phenylephrine HCl,
Doxylamine Succinate, Dextromethorphan HBr



3 23900 03846 2



ACTUAL SIZE

Headache, Fever, Sore Throat,
Minor Aches & Pains
Chest Congestion, Thins & Loosens Mucus
Nasal Congestion, Sinus Pressure
Cough

32 DAYQUIL COATED CAPLETS

Headache, Fever, Sore Throat,
Minor Aches & Pains
Nasal Congestion, Sinus Pressure
Sneezing, Runny Nose
Cough

16 NYQUIL COATED CAPLETS: COATED CAPLETS

DIST. BY PROCTER & GAMBLE,
CINCINNATI OH 45202
P&G www.pg.com
www.vicks.com
Patents: www.pg.com/patents
90884399

Drug Facts NyQuil + VapoCool Cold & Flu Active ingredients (in each caplet) Acetaminophen 325 mg Pain reliever/fever reducer Dextromethorphan HBr 10 mg Cough suppressant Doxylamine succinate 6.25 mg Antihistamine Phenylephrine HCl 5 mg Nasal decongestant Uses temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • cough to help you sleep • minor aches & pains • headache • fever • sore throat • runny nose & sneezing • reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take • more than 8 caplets in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product	Drug Facts DayQuil + VapoCool Cold & Flu Active ingredients (in each caplet) Acetaminophen 325 mg Pain reliever/fever reducer Dextromethorphan HBr 10 mg Cough suppressant Guaifenesin 200 mg Expectorant Phenylephrine HCl 5 mg Nasal decongestant Uses temporarily relieves common cold/flu symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • minor aches & pains • headache • fever • sore throat • reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage • helps loosen phlegm (mucus) and thin bronchial secretions to aid the bronchial passageways of bothersome mucus and make coughs more productive. Warnings Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take
--	---

Drug Facts (continued) Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema • trouble urinating due to enlarged prostate gland Ask a doctor or pharmacist before use if you are • taking sedatives or tranquilizers • taking the blood thinning drug warfarin When using this product • do not use more than directed • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.	Drug Facts (continued) • more than 8 caplets in 24 hours, which is the maximum daily amount for this product • with other drugs containing acetaminophen • 3 or more alcoholic drinks every day while using this product Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash If a skin reaction occurs, stop use and seek medical help right away Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly. Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. Ask a doctor before use if you have • liver disease • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin. When using this product, do not use more than directed. Stop use and ask a doctor if • you get nervous, dizzy or sleepless • pain, nasal congestion, or cough gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
---	--

DayQuil

POWERFUL RELIEF

NyQuil

DAY & NIGHT PACK

POWERFUL RELIEF

DayQuil

THE DAYTIME COUGHING, ACHING, STUFFY HEAD, FEVER, SORE THROAT, POWER THROUGH YOUR DAY, MEDICINE.

NyQuil

THE NIGHTTIME SNIFFLING, SNEEZING, COUGHING, ACHING, FEVER, BEST SLEEP WITH A COLD, MEDICINE.

Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Cough

Headache, Fever, Sore Throat, Minor Aches & Pains

Sneezing, Runny Nose

Cough

Non-Drowsy Alcohol Free Nighttime Relief Alcohol 10%

2 BOTTLES - 1 DAYQUIL/1 NYQUIL 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)

Parents: Learn about their children's medicine at www.StayWellOnYourSides.com

DIST. BY PROCTER & GAMBLE, CINCINNATI OH 45202

P&G
www.pg.com
www.nyquil.com
Patents: www.pg.com/patents

NyQuil Cold & Flu

Drug Facts	Active Ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg		Pain reliever/fever reducer
Dextromethorphen HBr 30 mg		Cough suppressant
Doxylamine succinate 12.5 mg		Antihistamine

Uses temporarily relieves common cold/flu symptoms:

- cough, due to minor throat and bronchial irritation
- headache
- stuffy nose and sneezing

Warnings

Do not use

Ask a doctor before use if you have

When using this product

Stop use and ask a doctor if

Directions

DayQuil Cold & Flu

Drug Facts	Active Ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg		Pain reliever/fever reducer
Dextromethorphen HBr 15 mg		Cough suppressant
Phenylephrine HCl 5 mg		Nasal decongestant

Uses temporarily relieves common cold/flu symptoms:

- cough, due to minor throat & bronchial irritation
- headache
- stuffy nose and sneezing

Warnings

Do not use

Ask a doctor before use if you have

When using this product, do not use more than directed.

Stop use and ask a doctor if

Directions

DayQuil

THE DAYTIME COUGHING, ACHING, STUFFY HEAD, FEVER, SORE THROAT. POWER THROUGH YOUR DAY, MEDICINE.

DayQuil

POWERFUL RELIEF

DAY & NIGHT PACK

NyQuil

THE NIGHTTIME, SNEEZING, SNEEZING, COUGHING, ACHING, FEVER. BEST SLEEP WITH A COLD, MEDICINE.

Parents:

P&G



<p>Drug Facts NyQuil™ Cold & Flu</p> <p>Active ingredients (in each LiquiCap)</p> <p>Acetaminophen 325 mg Pain reliever/fever reducer</p> <p>Dextromethorphan HBr 15 mg Cough suppressant</p> <p>Doxylamine succinate 6.25 mg Antihistamine</p> <p>Uses temporarily relieves common cold/flu symptoms:</p> <ul style="list-style-type: none"> cough due to minor throat & bronchial irritation headache minor aches & pains fever sore throat runny nose & sneezing <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> more than 8 LiquiCaps in 24 hours, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> skin reddening blisters rash 	<p>Drug Facts DayQuil™ Cold & Flu</p> <p>Active ingredients (in each LiquiCap)</p> <p>Acetaminophen 325 mg Pain reliever/fever reducer</p> <p>Dextromethorphan HBr 10 mg Cough suppressant</p> <p>Phenylephrine HCl 5 mg Nasal decongestant</p> <p>Uses temporarily relieves common cold/flu symptoms:</p> <ul style="list-style-type: none"> nasal congestion sore throat headache minor aches & pains fever cough due to minor throat & bronchial irritation <p>Warnings</p> <p>Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take</p> <ul style="list-style-type: none"> more than 8 LiquiCaps in 24 hours, which is the maximum daily amount for this product with other drugs containing acetaminophen 3 or more alcoholic drinks every day while using this product <p>Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:</p> <ul style="list-style-type: none"> skin reddening blisters rash 	<p>Drug Facts (continued)</p> <p>If a skin reaction occurs, stop use and seek medical help right away</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> liver disease glaucoma cough that occurs with too much phlegm (mucus) a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema trouble urinating due to enlarged prostate gland <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking sedatives or tranquilizers taking the blood thinning drug warfarin <p>When using this product</p> <ul style="list-style-type: none"> do not use more than directed excitability may occur, especially in children marked drowsiness may occur avoid alcoholic drinks be careful when driving a motor vehicle or operating machinery alcohol, sedatives, and tranquilizers may increase drowsiness <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> pain or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. <p>These could be signs of a serious condition.</p>	<p>Drug Facts (continued)</p> <p>If a skin reaction occurs, stop use and seek medical help right away</p> <p>Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.</p> <p>Do not use</p> <ul style="list-style-type: none"> with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> liver disease heart disease high blood pressure diabetes trouble urinating due to enlarged prostate gland cough that occurs with too much phlegm (mucus) persistent or chronic cough such as occurs with smoking, asthma, or emphysema <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking the blood thinning drug warfarin <p>When using this product, do not use more than directed.</p> <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> you get nervous, dizzy or sleepless pain, nasal congestion, or cough gets worse or lasts more than 7 days fever gets worse or lasts more than 3 days redness or swelling is present new symptoms occur cough comes back or occurs with rash or headache that lasts. <p>These could be signs of a serious condition.</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p>
<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <ul style="list-style-type: none"> take only as directed do not exceed 8 LiquiCaps per 24 hrs adults & children 12 yrs & over 2 LiquiCaps with water every 6 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use <p>Other information</p> <ul style="list-style-type: none"> store at no greater than 25°C <p>Inactive ingredients D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide</p> <p>Questions? 1-800-362-1683</p>	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <ul style="list-style-type: none"> take only as directed do not exceed 8 LiquiCaps per 24 hrs adults & children 12 yrs & over 2 LiquiCaps with water every 6 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use <p>Other information</p> <ul style="list-style-type: none"> store at no greater than 25°C <p>Inactive ingredients D&C Yellow No. 10, FD&C Blue No. 1, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide</p> <p>Questions? 1-800-362-1683</p>	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <ul style="list-style-type: none"> take only as directed do not exceed 8 LiquiCaps per 24 hrs adults & children 12 yrs & over 2 LiquiCaps with water every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use <p>Other information</p> <ul style="list-style-type: none"> store at no greater than 25°C <p>Inactive ingredients FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide</p> <p>Questions? 1-800-362-1683</p>	<p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.</p> <p>Directions</p> <ul style="list-style-type: none"> take only as directed do not exceed 8 LiquiCaps per 24 hrs adults & children 12 yrs & over 2 LiquiCaps with water every 4 hrs children 4 to under 12 yrs ask a doctor children under 4 yrs do not use <p>Other information</p> <ul style="list-style-type: none"> store at no greater than 25°C <p>Inactive ingredients FD&C Red No. 40, FD&C Yellow No. 6, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, sorbitol sorbitan solution, titanium dioxide</p> <p>Questions? 1-800-362-1683</p>

NyQuil SEVERE

Drug Facts

Active ingredients (in each 30 mL):

Ingredient	Purpose
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 15 mg	Sleep aid
Phenylephrine HCl 10 mg	Nasal decongestant

Uses: Temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily reduces throat breathing through the nose
- promotes nasal and/or sinus drainage

Warnings: Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- any other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use: If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) for depression, psychiatric or emotional condition, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a breathing problem or chronic cough that lasts or is associated with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are:

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product:

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- do not allow driving a motor vehicle or operating machinery, alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if:

- you get nervous, dizzy or sleepless
- you have congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

Directions:

- take only as directed
- only use the dose cap provided
- do not exceed 4 doses per 24 hrs
- adults & children 12 yrs & over 150 mL every 4 hrs
- children 6 to 11 yrs 100 mL every 4 hrs
- children 4 to 5 yrs 50 mL every 4 hrs
- children under 4 yrs ask a doctor
- do not use

Other information:

- each 30 mL contains: sodium 81 mg
- store at or below 75°F (23°C) and protect from light

Inactive ingredients: alcohol, citric acid, DSC Yellow No. 10, FD&C Green No. 3, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium bicarbonate, sodium chloride, sodium citrate, sorbitol, sucralose

Questions? 1-800-362-1883

DayQuil SEVERE

Drug Facts

Active ingredients (in each 15 mL):

Ingredient	Purpose
Dextromethorphan HBr 10 mg	Cough suppressant
Guafenesin 100 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses: Temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- minor aches & pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily reduces throat breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and the bronchial secretions to rid the bronchial passageways of mucus and make coughs more productive

Warnings: Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- any other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use: If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) for depression, psychiatric or emotional condition, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a breathing problem or chronic cough that lasts or is associated with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are:

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product:

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- do not allow driving a motor vehicle or operating machinery, alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if:

- you get nervous, dizzy or sleepless
- you have congestion, or cough gets worse or lasts more than 7 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

Directions:

- take only as directed
- only use the dose cap provided
- do not exceed 4 doses per 24 hrs
- adults & children 12 yrs & over 30 mL every 4 hrs
- children 6 to 11 yrs 15 mL every 4 hrs
- children 4 to 5 yrs 7.5 mL every 4 hrs
- children under 4 yrs ask a doctor
- do not use

Other information:

- each 15 mL contains: sodium 47 mg
- store at or below 75°F (23°C) and protect from light

Inactive ingredients: citric acid, DSC Yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose

Questions? 1-800-362-1883

NyQuil SEVERE

Drug Facts

Active ingredients (in each 30 mL):

Ingredient	Purpose
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 15 mg	Sleep aid
Phenylephrine HCl 10 mg	Nasal decongestant

Uses: Temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- minor aches & pains
- headache
- fever
- sore throat
- reduces swelling of nasal passages
- temporarily reduces throat breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and the bronchial secretions to rid the bronchial passageways of mucus and make coughs more productive

Warnings: Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4 doses in 24 hours, which is the maximum daily amount for this product
- any other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Do not use: If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If you are now taking a prescription monoamine oxidase inhibitor (MAOI) for depression, psychiatric or emotional condition, or Parkinson's disease, or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- a breathing problem or chronic cough that lasts or is associated with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are:

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

When using this product:

- do not use more than directed
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- do not allow driving a motor vehicle or operating machinery, alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if:

- you get nervous, dizzy or sleepless
- you have congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

Directions:

- take only as directed
- only use the dose cap provided
- do not exceed 4 doses per 24 hrs
- adults & children 12 yrs & over 150 mL every 4 hrs
- children 6 to 11 yrs 100 mL every 4 hrs
- children 4 to 5 yrs 50 mL every 4 hrs
- children under 4 yrs ask a doctor
- do not use

Other information:

- each 30 mL contains: sodium 81 mg
- store at or below 75°F (23°C) and protect from light

Inactive ingredients: alcohol, citric acid, DSC Yellow No. 10, FD&C Green No. 3, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium bicarbonate, sodium chloride, sodium citrate, sorbitol, sucralose

Questions? 1-800-362-1883

DayQuil SEVERE

THE DAYTIME, COUGHING, ACHING, STUFFY HEAD, FEVER, SORE THROAT, CONGESTED CHEST, POWER THROUGH YOUR DAY, MEDICINE.

Acetaminophen, Guafenesin, Phenylephrine HCl, Doxylamine Succinate, Pseudoephedrine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains

Chest Congestion, Thins & Loosens Mucus

Nasal Congestion, Sinus Pressure

Cough

Non-Drowsy Alcohol Free

NyQuil SEVERE

THE NIGHTTIME, SNEEZING, SNEEZING, COUGHING, ACHING, STUFFY HEAD, FEVER, BEST SLEEP WITH A COLD, MEDICINE.

Acetaminophen, Phenylephrine HCl, Doxylamine Succinate, Pseudoephedrine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Sneezing, Runny Nose

Cough

Nighttime Relief Alcohol 10%

PARENTS:
Learn more about us at www.StayMedicineRight.org

DIET: BY PROCTER & GAMBLE, CINCINNATI OH 45202

P&G
www.pg.com
Patents: www.pg.com/patents

2 BOTTLES - 1 DAYQUIL/1 NYQUIL 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)

DayQuil SEVERE

THE DAYTIME, COUGHING, ACHING, STUFFY HEAD, FEVER, SORE THROAT, CONGESTED CHEST, POWER THROUGH YOUR DAY, MEDICINE.

Acetaminophen, Guafenesin, Phenylephrine HCl, Doxylamine Succinate, Pseudoephedrine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains

Chest Congestion, Thins & Loosens Mucus

Nasal Congestion, Sinus Pressure

Cough

Non-Drowsy Alcohol Free

NyQuil SEVERE

THE NIGHTTIME, SNEEZING, SNEEZING, COUGHING, ACHING, STUFFY HEAD, FEVER, BEST SLEEP WITH A COLD, MEDICINE.

Acetaminophen, Phenylephrine HCl, Doxylamine Succinate, Pseudoephedrine HCl

Headache, Fever, Sore Throat, Minor Aches & Pains

Nasal Congestion, Sinus Pressure

Sneezing, Runny Nose

Cough



Nighttime Relief Alcohol 10%

PARENTS:
Learn more about us at www.StayMedicineRight.org

DIET: BY PROCTER & GAMBLE, CINCINNATI OH 45202

P&G
www.pg.com
Patents: www.pg.com/patents

2 BOTTLES - 1 DAYQUIL/1 NYQUIL 12 FL OZ (354 mL) EACH; TOTAL 24 FL OZ (708 mL)







Drug Facts (continued)

Uses temporarily relieves common cold/flu symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- sore throat
- minor aches & pains
- headache
- fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if:

- adult takes more than 4 doses (30 mL each) in 24 hrs, which is the maximum daily amount for this product
- child takes more than 4 doses (15 mL each) in 24 hrs, which is the maximum daily amount for this product
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day

Drug Facts (continued)

while using this product

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Drug Facts (continued)

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present

Drug Facts (continued)

- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	15 mL every 4 hrs
children 6 to under 12 yrs	ask a doctor
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 46 mg

Drug Facts (continued)

- store at no greater than 25°C and do not refrigerate

Inactive ingredients citric acid, FD&C yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum

Questions? 1-800-362-1883



www.pg.com
Patents: www.pg.com/patents



Drug Facts (continued)	
Warnings Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have <ul style="list-style-type: none">• heart disease• high blood pressure• thyroid disease• diabetes• trouble urinating due to enlarged prostate gland• cough that occurs with too much phlegm (mucus)• persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema• a sodium-restricted diet	
When using this product, do not use more than directed. Stop use and ask a doctor if <ul style="list-style-type: none">• you get nervous, dizzy or sleepless• symptoms do not improve within 7 days or occur with a fever• cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache These could be signs of a serious condition.	
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions <ul style="list-style-type: none">• take only as directed• only use the dose cup provided• do not exceed 4 doses per 24 hrs	
adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Drug Facts (continued)
Other information <ul style="list-style-type: none">• each 15 mL contains: sodium 45 mg• store at no greater than 25°C
Inactive ingredients citric acid, FD&C Yellow No. 6, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol, sucralose, xanthan gum
Questions? 1-800-362-1683



TRIPLE ACTION



DayQuil™

HOT REMEDY

RAPID RELIEF MULTI-SYMPTOM

**COLD & FLU
+ CONGESTION**

Acetaminophen,
Dextromethorphan HBr,
Phenylephrine HCl



Headache, Fever,
Sore Throat,
Minor Aches
& Pains

Nasal Congestion,
Sinus Pressure
Cough

P&G
www.pg.com
vicks.com
Patents: www.pg.com/patents

Non-Drowsy

**HONEY
LEMON
FLAVORED**

8 POWDER PACKETS

Tamper Evident: Do not use if sachet is opened or torn

Keep outer package for complete product information.

Drug Facts

Active Ingredients (in each Packet)	Purpose
Acetaminophen 500 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Phenylephrine HCl 10 mg	Nasal decongestant

Uses: temporarily relieves common cold/flu symptoms:

- minor aches & pains
- sore throat
- fever
- headache
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- nasal congestion
- sinus congestion & pressure
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warnings: This product contains acetaminophen. Severe liver damage may occur if you take more than 6 doses in 24 hours, which is the maximum daily amount for this product.

Allergy Alert: Acetaminophen may cause severe skin reactions. Symptoms may include: • Skin reddening • Blisters • Rash

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- If you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- liver disease • heart disease • high blood pressure • thyroid disease • diabetes
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product, do not use more than directed

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be signs of a serious condition.

Directions

- take only as directed
- do not exceed 6 doses per 24 hrs
- Dissolve contents of one packet into 8 oz. of hot water and stir briskly; sip while hot. Consume entire drink within 10-15 minutes
- If using a microwave, add contents of one packet to 8 oz. of cool water, stir briskly before and after heating. Do not overheat.

Age	Dose
adults & children 12 yrs & over	one packet (dose) every 4 hours
children under 12 yrs	do not use

Other information

- each packet contains: potassium 10 mg • phenylephrine: contains phenylephrine 64 mg per dose
- do not exceed 25°C.

Inactive ingredients: acetaminophen, potassium, aspartame, carbonylmethylcellulose, citric acid, FD&C Yellow No. 10, FD&C Blue No. 1, FD&C Red No. 40, flavors, sucrose, triethyl citrate, sodium phosphate

Questions?
1-800-368-1888

MADE IN MEXICO
DIST. BY: PROCTER & GAMBLE,
CINCINNATI, OH 45202
90284422

Drug Facts (continued)
 contact a Poison Control Center right away.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 15 mg
- store at no greater than 25°C

Inactive ingredients citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions? 1-800-362-1683

Drug Facts (continued)
 to rid the bronchial passageways of bothersome mucus and make coughs more productive.

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

When using this product, do not use more than directed.

Stop use and ask a doctor if:

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or

TAMPER EVIDENT:
 DO NOT USE IF PRINTED SHRINKBAND ON BOTTLE IS BROKEN OR MISSING.
 PEEL BACK FOR DRUG FACTS

Drug Facts

Active ingredients (in each 15 mL)

Active ingredients (in each 15 mL)	Purpose
Dextromethorphan HBr 10 mg	Cough suppressant
Guafenesin 100 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves common cold symptoms:
- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passage
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions

PARENTS:
 Learn about teen medicine abuse
www.StopMedicineAbuse.org

DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202
 Patents: www.pg.com/patents
vicks.com

P&G
www.pg.com
 90461664

Depend Bank
 Recycled Cap
 100% POST CONSUMER WASTE
 PLASTIC BOTTLE

2390003077 0

VICKS

DayQuil Kids

BERRY

COLD & COUGH + MUCUS

Guafenesin, Phenylephrine HCl, Dextromethorphan HBr

Chest Congestion
 Nasal Congestion
 Cough

AGES 6+

8 FL OZ (236 mL)

Non-Drowsy
 FREE OF ALCOHOL & ACETAMINOPHEN

90474161



Drug Facts (continued)

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- each 15 mL contains: sodium 14 mg
- store at no greater than 25°C

Inactive ingredients citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor (with honey), glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions? 1-800-362-1683

Drug Facts (continued)

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

When using this product, do not use more than directed.

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.



NyQuil Kids Honey COLD & COUGH + CONGESTION

Drug Facts

Active ingredients (in each 15 mL)

Ingredient	Purpose
Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine succinate 6.25 mg	Sleep aid
Phenylephrine HCl 5 mg	Nasal decongestant

Uses Temporarily relieves common cold symptoms:

- nasal congestion
- cough due to minor throat & bronchial irritation
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or occurs with smoking, asthma or emphysema

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product:

- do not use more than directed.
- excitability may occur, especially in children.
- marked drowsiness may occur.
- alcohol, sedatives, and tranquilizers may increase drowsiness.
- avoid alcoholic drinks.
- be careful when driving a motor vehicle or operating machinery.

Stop use and ask a doctor if:

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs
- adults & children 12 yrs & over 30 mL every 4 hrs
- children 6 to under 12 yrs 15 mL every 4 hrs
- children 4 to under 6 yrs do not use unless directed by a doctor
- children under 4 yrs do not use

Other information

- each 15 mL contains sodium 14 mg
- store at no greater than 20°C

Inactive ingredients citric acid, DAC Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor (with honey), glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions? 1-800-362-1883

DayQuil Kids Honey COLD & COUGH + MUCUS

Drug Facts

Active ingredients (in each 15 mL)

Ingredient	Purpose
Dextromethorphan HBr 10 mg	Cough suppressant
Sudafed 120 mg	Expectorant
Phenylephrine HCl 5 mg	Nasal decongestant

Uses Temporarily relieves common cold symptoms:

- nasal congestion
- sinus congestion & pressure
- cough due to minor throat & bronchial irritation
- cough to help you sleep
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passages of bothersome mucus and make coughs more productive.

Warnings

Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have:

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema

When using this product, do not use more than directed.

Stop use and ask a doctor if:

- you get nervous, dizzy or sleepless
- symptoms do not improve within 7 days or occur with a fever
- cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take only as directed
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs
- adults & children 12 yrs & over 30 mL every 4 hrs
- children 6 to under 12 yrs 15 mL every 4 hrs
- children 4 to under 6 yrs ask a doctor
- children under 4 yrs do not use

Other information

- each 15 mL contains sodium 14 mg
- store at no greater than 20°C

Inactive ingredients citric acid, D&C Yellow No. 10, FD&C Green No. 3, FD&C Red No. 40, FD&C Yellow No. 6, flavor (with honey), glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum

Questions? 1-800-362-1883

DayQuil Kids HONEY COLD & COUGH + MUCUS

Safe, fast and effective cough & cold relief for your little one.

Specially formulated for kids and flavored with **real honey** to coat and soothe their throat.

From **VICK'S®** the world's #1 selling cough & cold brand.

PLASTIC BOTTLE

TAMPER EVIDENT: Do not use if printed shrinkband seal around the neck is broken or missing

90402755

DayQuil Kids NyQuil Kids

VALUE COMBO PACK 1 DAY + 1 NIGHT

coats & soothes (flavored with real honey)

AGES 6+

Non-Drowsy

Free of Alcohol & Acetaminophen

2 BOTTLES - 1 DAYQUIL / 1 NYQUIL 8 FL OZ (236 mL) EACH; TOTAL 16 FL OZ (473 mL)

Ingredients:

- DayQuil: Gaisfensan, Phenylephrine HCl, Dextromethorphan HBr
- NyQuil: Doxylamine Succinate, Phenylephrine HCl, Dextromethorphan HBr

Relieves:

- Chest Congestion
- Nasal Congestion
- Cough
- Sneezing, Runny Nose
- Nasal Congestion
- Cough

NyQuil Kids HONEY COLD & COUGH + CONGESTION

Honey-luscious day and night cold, cough & congestion relief for your child.

DayQuil Relieves coughs, stuffy noses, and chest congestion for kids with a non-drowsy formula.

NyQuil Provides kids multi-symptom relief plus the best sleep with a cold medicine.

PARENTS: Learn about teen medicine abuse! www.StopMedicineAbuse.org

P&G www.pg.com vicks.com Patents: www.pg.com/patents

NyQuil™ Kids BERRY

Drug Facts	Purpose
Active ingredients (in each 15 mL) Chlorpheniramine maleate 2 mg Dextromethorphan HBr 15 mg	Antihistamine Cough suppressant
Uses Temporarily relieves common cold symptoms: • cough due to minor throat & bronchial irritation • sneezing • runny nose	
Warnings Do not use • if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. • to make a child sleepy	
Ask a doctor before use if you have • glaucoma • cough that occurs with too much phlegm (mucus) • a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis or emphysema • trouble urinating due to enlarged prostate gland	
Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers	
When using this product • excitability may occur, especially in children • marked drowsiness may occur • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • alcohol, sedatives, and tranquilizers may increase drowsiness	
Stop use and ask a doctor if • cough persists for more than 1 week, tends to recur, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition.	
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions • take only as directed • only use the dose cap provided • do not exceed 4 doses per 24 hrs	
adults & children 12 yrs & over 30 mL every 6 hrs children 6 to under 12 yrs 15 mL every 6 hrs children 4 to under 6 yrs do not use unless directed by a doctor children under 4 yrs do not use	
Other information • each 15 mL contains: sodium 13 mg • store at or greater than 25°C	
Inactive ingredients citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum	
Questions? 1-800-362-1683	

DayQuil™ Kids BERRY

Drug Facts	Purpose
Active ingredients (in each 15 mL) Dextromethorphan HBr 10 mg Guaifenesin 100 mg Phenylephrine HCl 5 mg	Cough suppressant Expectorant Nasal decongestant
Uses Temporarily relieves common cold symptoms: • nasal congestion • sinus congestion & pressure • cough due to minor throat & bronchial irritation • cough to help you sleep • reduces swelling of nasal passages • temporarily restores freer breathing through the nose • promotes nasal and/or sinus drainage • helps loosen phlegm (mucus) and this bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive.	
Warnings Do not use if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.	
Ask a doctor before use if you have • heart disease • high blood pressure • thyroid disease • diabetes • trouble urinating due to enlarged prostate gland • cough that occurs with too much phlegm (mucus) • persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema	
When using this product, do not use more than directed.	
Stop use and ask a doctor if • you get nervous, dizzy or sleepless • symptoms do not improve within 7 days or occur with a fever • cough persists for more than 7 days, comes back or occurs with a fever, rash or persistent headache. These could be signs of a serious condition.	
If pregnant or breast-feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions • take only as directed • only use the dose cap provided • do not exceed 4 doses per 24 hrs	
adults & children 12 yrs & over 30 mL every 4 hrs children 6 to under 12 yrs 15 mL every 4 hrs children 4 to under 6 yrs ask a doctor children under 4 yrs do not use	
Other information • each 15 mL contains: sodium 15 mg • store at or greater than 25°C	
Inactive ingredients citric acid, FD&C Red No. 40, flavor, glycerin, propylene glycol, saccharin sodium, sodium benzoate, sodium citrate, sorbitol, sucralose, water, xanthan gum	
Questions? 1-800-362-1683	

DayQuil Kids

BERRY
COLD & COUGH
+ MUCUS

Fast & effective
cough & cold relief
for your little one.

Specially formulated
for kids to treat their
most bothersome
cold symptoms.

From **VICKS®**
the world's #1 selling
cough & cold brand.

DayQuil Kids

NyQuil Kids

VALUE COMBO PACK
1 DAY + 1 NIGHT

NyQuil Kids

BERRY
COLD & COUGH
+ RUNNY NOSE

Berry-licious
day & night cold
& cough relief
for your child.

DayQuil™
relieves coughs,
stuffy noses, and
chest congestion
for kids with a
non-drowsy formula.

NyQuil™
provides kids
multi-symptom relief
plus the best sleep
with a cold medicine.

AGES 6+

Guafenesin,
Phenylephrine HCl,
Dextromethorphan HBr

Chest Congestion
Nasal Congestion
Cough

Non-Drowsy

FREE OF ALCOHOL & ACETAMINOPHEN

AGES 6+

Chlorpheniramine maleate,
Dextromethorphan HBr

Sneezing,
Runny Nose
Cough

Nighttime Relief

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org

P&G
www.pg.com
vicks.com
Patents: www.pg.com/patents

TAMPER EVIDENT. DO NOT USE IF
PRINTED SHRINK-BAND SEAL AROUND
THE NECK IS BROKEN OR MISSING

2 BOTTLES - 1 DAYQUIL / 1 NYQUIL 8 FL OZ (236 mL) EACH; TOTAL 16 FL OZ (472 mL)

239003101

90463277